

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- asthma
- taken a diuretic
- reached age 60 or older
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- taking any other drug containing an NSAID (prescription or nonprescription)
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking a blood thinning (anticoagulant) or steroid drug
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Other information

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- **do not take more than directed**
- do not take longer than 10 days, unless directed by a doctor (see WARNINGS)
- adults and children 12 years and over: take 2 caplets at bedtime.
- do not take more than 2 caplets in 24 hours
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Do Not Use if foil seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Manufactured for: **Dr. Reddy's Laboratories Limited** Bachepalli – 502 325 INDIA.

Inactive ingredients

carnauba wax, corn starch, colloidal silicon dioxide, croscarmellose sodium, FD&C blue no. 2, hypromellose, microcrystalline cellulose, polydextrose, polyethylene glycol 400, pregelatinized starch, sodium lauryl sulfate, stearic acid, titanium dioxide